fragile XE syndrome

Fragile XE syndrome is a genetic disorder that impairs thinking ability and cognitive functioning. Most affected individuals have mild intellectual disability. In some people with this condition, cognitive function is described as borderline, which means that it is below average but not low enough to be classified as an intellectual disability. Females are rarely diagnosed with fragile XE syndrome, likely because the signs and symptoms are so mild that the individuals function normally.

Learning disabilities are the most common sign of impaired cognitive function in people with fragile XE syndrome. The learning disabilities are likely a result of communication and behavioral problems, including delayed speech, poor writing skills, hyperactivity, and a short attention span. Some affected individuals display autistic behaviors, such as hand flapping, repetitive behaviors, and intense interest in a particular subject. Unlike some other forms of intellectual disability, cognitive functioning remains steady and does not decline with age in fragile XE syndrome.

Frequency

Fragile XE syndrome is estimated to affect 1 in 25,000 to 100,000 newborn males. Only a small number of affected females have been described in the medical literature. Because mildly affected individuals may never be diagnosed, it is thought that the condition may be more common than reported.

Genetic Changes

Fragile XE syndrome is caused by mutations in the *AFF2* gene. This gene provides instructions for making a protein whose function is not well understood. Some studies show that the AFF2 protein can attach (bind) to DNA and help control the activity of other genes. Other studies suggest that the AFF2 protein is involved in the process by which the blueprint for making proteins is cut and rearranged to produce different versions of the protein (alternative splicing). Researchers are working to determine which genes and proteins are affected by AFF2.

Nearly all cases of fragile XE syndrome occur when a region of the *AFF2* gene, known as the CCG trinucleotide repeat, is abnormally expanded. Normally, this segment of three DNA building blocks (nucleotides) is repeated approximately 4 to 40 times. However, in people with fragile XE syndrome, the CCG segment is repeated more than 200 times, which makes this region of the gene unstable. (When expanded, this region is known as the FRAXE fragile site.) As a result, the *AFF2* gene is turned off (silenced), and no AFF2 protein is produced. It is unclear how a shortage of this protein leads to intellectual disability in people with fragile XE syndrome.

People with 50 to 200 CCG repeats are said to have an *AFF2* gene premutation. Current research suggests that people with a premutation do not have associated cognitive problems.

Inheritance Pattern

Fragile XE syndrome is inherited in an X-linked dominant pattern. A condition is considered X-linked if the mutated gene that causes the disorder is located on the X chromosome, which is one of the two sex chromosomes. In females (who have two X chromosomes), a mutation in one of the two copies of the gene in each cell is sufficient to cause the disorder. In males (who have only one X chromosome), a mutation in the only copy of the gene in each cell causes the disorder. In most cases, males experience more severe symptoms of the disorder than females.

In parents with the *AFF*2 gene premutation, the number of CCG repeats can expand to more than 200 in cells that develop into eggs or sperm. This means that parents with the premutation have an increased risk of having a child with fragile XE syndrome. A characteristic of X-linked inheritance is that fathers cannot pass X-linked traits to their sons; sons receive a Y chromosome from their father, which does not include the *AFF*2 gene.

Other Names for This Condition

- FRAXE intellectual deficit
- FRAXE intellectual disability
- FRAXE mental retardation syndrome
- FRAXE syndrome
- mental retardation, X-linked, associated with fragile site FRAXE
- mental retardation, X-linked, FRAXE type

Diagnosis & Management

Genetic Testing

 Genetic Testing Registry: FRAXE https://www.ncbi.nlm.nih.gov/gtr/conditions/C0751157/

Other Diagnosis and Management Resources

 Centers for Disease Control and Prevention: Developmental Screening Fact Sheet https://www.cdc.gov/ncbddd/actearly/pdf/parents_pdfs/ DevelopmentalScreening.pdf

General Information from MedlinePlus

- Diagnostic Tests
 https://medlineplus.gov/diagnostictests.html
- Drug Therapy https://medlineplus.gov/drugtherapy.html
- Genetic Counseling https://medlineplus.gov/geneticcounseling.html
- Palliative Care https://medlineplus.gov/palliativecare.html
- Surgery and Rehabilitation https://medlineplus.gov/surgeryandrehabilitation.html

Additional Information & Resources

MedlinePlus

- Encyclopedia: Intellectual Disability https://medlineplus.gov/ency/article/001523.htm
- Health Topic: Developmental Disabilities
 https://medlineplus.gov/developmentaldisabilities.html

Genetic and Rare Diseases Information Center

 Fragile XE syndrome https://rarediseases.info.nih.gov/diseases/2378/fragile-xe-syndrome

Additional NIH Resources

 Eunice Kennedy Shriver National Institute of Child Health and Human Development: Intellectual and Developmental Disabilities https://www.nichd.nih.gov/health/topics/idds/conditioninfo/Pages/default.aspx

Educational Resources

- Centers for Disease Control and Prevention: Intellectual Disability https://www.cdc.gov/ncbddd/actearly/pdf/parents_pdfs/IntellectualDisability.pdf
- Disease InfoSearch: Fragile XE Syndrome http://www.diseaseinfosearch.org/Fragile+XE+Syndrome/2911
- MalaCards: mental retardation, x-linked, fraxe type
 http://www.malacards.org/card/mental_retardation_x_linked_fraxe_type

- My 46 Trait Profile https://www.my46.org/trait-document?trait=Fragile%20XE%20syndrome&parent=Genetic%20Syndromes&type=profile
- Orphanet: FRAXE intellectual disability http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=100973

Patient Support and Advocacy Resources

- Resource List from the University of Kansas Medical Center: Developmental Delay http://www.kumc.edu/gec/support/devdelay.html
- The Arc: For People With Intellectual and Developmental Disabilities http://www.thearc.org/

Scientific Articles on PubMed

PubMed

https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28fraxe+intellectual+disability %5BTIAB%5D%29+OR+%28fraxe+syndrome%5BTIAB%5D%29+OR+%28FRAXE %5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D

OMIM

 MENTAL RETARDATION, X-LINKED, ASSOCIATED WITH FRAGILE SITE FRAXE http://omim.org/entry/309548

Sources for This Summary

- Gecz J, Gedeon AK, Sutherland GR, Mulley JC. Identification of the gene FMR2, associated with FRAXE mental retardation. Nat Genet. 1996 May;13(1):105-8.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/8673085
- Gecz J. The FMR2 gene, FRAXE and non-specific X-linked mental retardation: clinical and molecular aspects. Ann Hum Genet. 2000 Mar;64(Pt 2):95-106. Review.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/11246464
- Knight SJ, Flannery AV, Hirst MC, Campbell L, Christodoulou Z, Phelps SR, Pointon J, Middleton-Price HR, Barnicoat A, Pembrey ME, et al. Trinucleotide repeat amplification and hypermethylation of a CpG island in FRAXE mental retardation. Cell. 1993 Jul 16;74(1):127-34.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/8334699
- Melko M, Douguet D, Bensaid M, Zongaro S, Verheggen C, Gecz J, Bardoni B. Functional characterization of the AFF (AF4/FMR2) family of RNA-binding proteins: insights into the molecular pathology of FRAXE intellectual disability. Hum Mol Genet. 2011 May 15;20(10):1873-85. doi: 10.1093/hmg/ddr069. Epub 2011 Feb 17.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/21330300

Reprinted from Genetics Home Reference:

https://ghr.nlm.nih.gov/condition/fragile-xe-syndrome

Reviewed: January 2014 Published: March 21, 2017

Lister Hill National Center for Biomedical Communications U.S. National Library of Medicine National Institutes of Health Department of Health & Human Services